

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 214 879 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**19.06.2002 Bulletin 2002/25**

(21) Application number: **01403208.0**

(22) Date of filing: **12.12.2001**

(51) Int Cl.7:

**A01N 35/04**, A01N 35/02,  
A01N 43/08, A01N 31/02,  
A01N 31/04, A01N 31/06,  
A01N 31/08, A01N 37/40,  
A61K 7/46  
// (A01N35/04, 43:08, 37:40,  
35:02, 31:08, 31:06, 31:04,  
31:02),  
(A01N35/02, 43:08, 37:40,  
35:04, 31:08, 31:06, 31:04,  
31:02)

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **12.12.2000 JP 2000376958**

(71) Applicant: **Takasago International Corporation  
Tokyo 144-8721 (JP)**

(72) Inventors:

- **Hanada, Minoru**  
**5-chome, Ohta-ku, Tokyo 144-8721 (JP)**

• **Iwai, Hisao**

**Kamata 5 chome, Ohta-ku, Tokyo 144-8721 (JP)**

• **Fujigasaki, Jun**

**Kamata 5-chome, Ohta-ku, Tokyo 144-8721 (JP)**

(74) Representative: **Uchida, Kenji et al**

**S.A. Fedit-Loriot et Autres Conseils en Propriété  
Industrielle,  
38, avenue Hoche  
75008 Paris (FR)**

(54) **Antifungal fragrance composition**

(57) An antifungal fragrance composition containing as its active ingredients: (a) at least one type of fragrance component selected from the group consisting of an aliphatic or aromatic aldehyde having 5 to 16 carbon atoms, an aliphatic alcohol having 5 to 16 carbon atoms and a phenol, and (b) at least one type of compound selected from the group consisting of an aliphatic

or aromatic aldehyde, aliphatic alcohol, phenol, acetal and ester demonstrating synergistic effects in the presence of said fragrance component. The present antifungal fragrance composition demonstrates antifungal activity by adding in a small amount that only has a minimal effect on the scent of the fragrance component.

**EP 1 214 879 A2**

## Description

## BACKGROUND OF THE INVENTION

## 5 Field of the Invention

[0001] The present invention relates to a fragrance composition having antifungal activity, and the object of the present invention is to provide a highly safe antifungal fragrance composition that disperses an arbitrary fragrance in the room, closet or shoe box in which it is placed while simultaneously inhibiting mold growth, thereby preventing  
10 contamination of wall surfaces, floor surfaces, shoe surfaces and so forth caused by mold.

## Prior Art

[0002] One of the characteristics of harmful mold growth in homes in Japan is the high temperature and humidity  
15 that occur in summer. In contrast, during the winter, in the case of concrete housing complexes such as apartments in particular, a major cause of harmful mold growth is the interior condensation of moisture due to thermal insulation work being performed by internal thermal insulating methods. For these reasons, a large amount of mold is detected indoors in both the summer and winter. The types of mold isolated from rooms and bathrooms have been reported to vary according to the season, the region in which they were researched and other factors (Antibacterial and Antifungal  
20 Agents, Vol. 19, pp. 127-134 (1991); Antibacterial and Antifungal Agents, Vol. 28, pp. 421-426 (2000); etc.).

[0003] For example, *Wallemia*, *Aspergillus*, *Cladosporium*, *Penicillium* and other species of molds have been isolated from floor dust, while mold in the form of black stains present in the joints of bathroom tiles are said to be frequently caused by mold species such as *Cladosporium*, *Phoma*, *Auerobasidium* and *Ochroconis*. When such molds grow on  
25 wall surfaces, in closets or in shoe boxes, they can cause numerous problems such as mildew and other unpleasant odors, the creation of white, yellow, black or other stains, damage to wall surfaces or the resulting spores can be become suspended in the room and cause allergies.

[0004] Antifungal agents are frequently used as a means of preventing such mold growth. However, many of the antifungal agents that are commonly used are aqueous sprays containing sodium hypochlorite as their main ingredient. Although these antifungal agents are easy to use since they can be rinsed off after use if used in locations involving  
30 the use of water such as in bathrooms or lavatories, in the case of ordinary rooms, closets and shoe boxes, etc., there are concerns over their strongly irritating odor and safety with respect to the human body. Moreover, there is also the risk of the moisture remaining after spraying becoming a hotbed for mold growth in the form of new moisture resulting following decomposition of the active ingredient such as sodium hypochlorite.

[0005] Essential oils and their components, fragrant chemical products, have been widely known to have antibacterial and antifungal activity since long ago (Toru Asagoshi, Journal of the Society of Cosmetic Chemists of Japan, Vol. 34,  
35 pp. 25-46 (2000) and cited references therein), and have been shown in numerous reports to demonstrate activity not only in methods involving direct contact with test microorganisms, but also in the vapor state (Maruzzella, J.C. et al., American Perfumer and Aromatics 74 (Aug), 21-22 (1959); Shinobu Gocho, Antibacterial and Antifungal Agents, Vol. 20, p. 585 (1992); Tamio Nishimura, Aromatopia, 10, 60-63 (1995)).

[0006] Examples of patents describing the demonstration of antibacterial or antifungal activity by essential oils and fragrant chemical products in the vapor state include a non-therapeutic antimicrobial (Japanese Patent Publication No. Hei 3-77161), a bacteriostatic aromatic agent for the bathroom (Japanese Unexamined Patent Publication No. Hei 6-24952), an antimicrobial activity detection method (Japanese Unexamined Patent Publication No. Hei 10-108691), a fungal infection and diffusion preventive agent using essential oil (Japanese Unexamined Patent Publication No. Hei  
45 10-338630), a antibacterial and bactericidal agent for food (Japanese Unexamined Patent Publication No. Hei 11-332534) and a bactericidal agent for the refrigerator (Japanese Unexamined Patent Publication No. Hei 11-335219).

[0007] However, the antibacterial and antifungal activities of these fragrant components are not so potent, and when prepared in an amount that demonstrates activity, the characteristic aroma of the active ingredient becomes conspicuous, and there are many cases in which the scent of the overall preparation is controlled by the aroma of the active  
50 ingredient, thereby making it difficult to generate antibacterial and antifungal activity while maintaining a specific scent (such as a citrus, fruity, woody or mint scent).

## SUMMARY OF THE INVENTION

[0008] Therefore, the object of the present invention is to provide an antifungal fragrance composition that is able to demonstrate antifungal activity with addition of a small amount having a minimal effect on the scent of the fragrant component.

[0009] As a result of conducting earnest research on the antibacterial activity and antifungal activity of fragrant vapor,

the inventors of the present invention found that by mixing at least one type of fragrance component selected from the group consisting of aliphatic or aromatic aldehydes having 5 to 16 carbon atoms, aliphatic alcohols having 5 to 16 carbon atoms and phenol derivatives, and at least one type of compound selected from the group consisting of aliphatic or aromatic aldehydes, aliphatic alcohols, phenol derivatives, acetals and esters, its activity is synergistically enhanced and antifungal activity can be demonstrated even when both components are mixed at low concentrations such as those at which activity is not demonstrated with the fragrance component alone, and consequently not disturbing its specific scent, and thus they accomplished the present invention.

**[0010]** Namely, the present invention is:

1) an antifungal fragrance composition containing as its active ingredients: (a) at least one type of fragrance component selected from the group consisting of an aliphatic or aromatic aldehyde having 5 to 16 carbon atoms, an aliphatic alcohol having 5 to 16 carbon atoms and a phenol, and (b) at least one type of compound selected from the group consisting of an aliphatic or aromatic aldehyde, aliphatic alcohol, phenol, acetal and ester demonstrating synergistic effects in the presence of the above-mentioned fragrance component;

2) the antifungal fragrance composition described in 1) above, wherein the fragrance composition of the component (a) is selected from the group consisting of citral, citronellal, trans-2-hexenal, octyl aldehyde, trans-2-undecenal, 10-undecen-1-al, dimethyl tetrahydrobenzaldehyde, benzaldehyde, salicylic aldehyde, 3-phenylpropanal, cuminaldehyde, furfural, geraniol, citronellol, 9-decen-1-ol, 1-decanol and eugenol;

3) the antifungal fragrance composition described in 1) or 2) above, wherein the compound demonstrating synergistic effects of the component (b) is citral, citronellal, trans-2-hexenal, octylaldehyde, 1-nonanal, trans-2-undecenal, undecylenealdehyde, dimethyltetrahydrobenzaldehyde, benzaldehyde, salicylic aldehyde, 3-phenylpropionic aldehyde, cuminaldehyde, furfural, phenylacetaldehyde, anisaldehyde, geraniol, citronellol, 9-decenol, 1-decanol, thymol, eugenol, citral dimethylacetal (hereinafter abbreviated as citral DMA), citral diethylacetal (hereinafter abbreviated as citral DEA), cinnamic aldehydodimethylacetal (hereinafter abbreviated as cinnamic aldehyde DMA), or methyl salicylate;

4) the antifungal fragrance composition according to any one of 1) to 3) above that is contained at 1 to 50% by weight in a gel;

5) the antifungal fragrance composition according to any one of 1) to 3) above that is contained at 1 to 60% by weight in a powder or granules;

6) the antifungal fragrance composition according to any one of 1) to 3) above that is contained at 1 to 30% by weight in a liquid;

7) an antifungal fragrance composition package comprising the containing of the antifungal fragrance composition described in any one of 1) to 3) above, or the composition dissolved in a volatile solvent, in a gas permeable container;

8) the antifungal fragrance composition package described in 7) above, wherein the gas permeable container is a pouch-shaped object made of a laminate comprising a gas permeable thermoplastic resin on the inside and a gas permeable support on the outside;

9) the antifungal fragrance composition package described in 8) above, wherein the gas permeable support is a non-woven fabric;

10) an antifungal fragrance composition package comprising the containing of the antifungal fragrance composition described in any one of 1) to 3) above, or the composition dissolved in a volatile solvent, and a propellant in a spray can; and,

11) a method of inhibiting mold growth comprising: placing the antifungal fragrance composition package described in any one of 7) to 9) above at an arbitrary location, and dispersing the components (a) and (b) in the antifungal fragrance composition contained in the package.

**[0011]** In the present invention, by combining the use of the fragrant component of the component (a) having antifungal activity with the component (b) demonstrating a synergistic effect in the presence of the component (a), the

amount of fragrant component blended that demonstrates activity can be reduced to a much lower amount than in the case of using the component (a) alone. As a result, even in the case of blending into other fragrances, there are minimal effects on the fragrant component, thereby making it possible to provide an antifungal fragrance composition that maintains an arbitrary aroma while exhibiting an extremely high degree of safety with respect to the human body.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

**[0012]** Examples of aliphatic or aromatic aldehydes that can be used in the present invention include citral, citronellal, trans-2-hexenal, octylaldehyde, trans-2-undecenal, undecylaldehyde (10-undecenal or undecenoic aldehyde), dimethyltetrahydrobenzaldehyde (tripural), benzaldehyde, salicylaldehyde, 3-phenylpropanal, cuminaldehyde and furfural, as well as nonylaldehyde, phenylacetaldehyde, cinnamic aldehyde, anisaldehyde, perillaldehyde and helional. In addition, acetals of these aldehydes such as dimethylacetals and diethylacetals can similarly also be used. Moreover, in addition to these fragrant chemical products, natural fragrances and essential oils containing these substances as their components can also be used.

**[0013]** Examples of alcohols and phenols used in the present invention include geraniol, citronellol, 9-decenol, decanol and eugenol as well as thymol, isoeugenol, hinokitiol, perilla alcohol. Moreover, natural fragrances and essential oils containing these substances as their components can also be used.

**[0014]** Examples of acetals that can be used include citral DMA, citral DEA and cinnamic aldehyde DMA, while examples of esters that can be used include methyl salicylate.

**[0015]** Although the blending ratio of the component (a) and component (b) in the antifungal fragrance composition of the present invention differs according to the type of each component and cannot be unconditionally determined, normally a ratio of (a):(b) = 1:0.01 to 50 is preferable. For example, in the case of component (a) being citral and component (b) being citronellal, about 1 to 20 parts by weight of citronellal should be blended relative to 1 part by weight of citral.

**[0016]** The antifungal fragrance composition of the present invention can be in the form of a gel, powder, granules, liquid and so forth. In the case of being in the form of a gel, although it is common to use a gelling agent, there are no restrictions on the gelling agent used, with examples of gelling agents that can be used including those formed from water-soluble high polymeric organic substances such as polyvinyl alcohol, polyethylene glycol, polyvinyl pyrrolidone, alginic acid, cellulose, starch, their derivatives or their salts, synthetic resin emulsions such as vinyl acetate resin emulsion and copolymer-based resin emulsions, e.g. those of vinyl acetate and vinyl chloride, aqueous emulsions such as rubber-based latex, gelling agents derived from naturally-occurring substances extracted from various animals, plants, algae and microorganisms such as carrageenan, agar, gelatin and pectin. Surfactants such as sodium stearate and sodium isostearate may be used to disperse and emulsify the antifungal fragrance component in the gel. In the case of being in the form of a gel, the gel may contain 1 to 50% by weight of the antifungal fragrance composition.

**[0017]** In addition, in the case of being in the form of a powder or granules, 1 to 60% by weight of the antifungal fragrance composition should be adsorbed or impregnated. Examples of the carrier of the powder or granules include silica gel, zeolite, calcium silicate, diatomaceous earth, activated charcoal, alumina, allophane and vermiculite.

**[0018]** In addition, the antifungal fragrance composition of the present invention may also be a liquid, and in this case, can be composed of a surfactant such as polyoxyethylene hardened castor oil or 1 to 40% by weight aqueous ethanol as the solubilizing agent of the antifungal fragrance component, antioxidant such as or dibutylhydroxytoluene, ultraviolet absorber such as urocanic acid or oxybenzone and a pigment, etc., and the antifungal fragrance component should be blended at 1 to 30% by weight.

**[0019]** The present invention also provides an antifungal fragrance composition package that contains the above antifungal fragrance composition in a container. The antifungal fragrance composition is contained directly, dissolved in a volatile solvent such as alcohol, directly as a liquid, or in a gas permeable container when the composition is in the form of the above-mentioned gel, powder or granules. Examples of gas permeable containers that are used preferably include those in which laminated paper is formed into the shape of a pouch, the laminated paper being made by using a non-woven fabric composed of a gas permeable material for the support, and laminating onto the support a film having an inner surface of polyethylene film or other gas permeable, thermoplastic film.

**[0020]** In addition, the present invention can also provide an antifungal fragrance composition package in which the antifungal fragrance composition of the present invention is placed in a manual spray container either directly or after dissolving in a volatile solvent such as alcohol, or charged into a spray can with a propellant such as LPG, dimethyl ether or liquefied carbon dioxide gas.

**[0021]** If the above antifungal fragrance composition package is placed in a location such as a room, closet or shoe box, together with the scent of the fragrance component being dispersed at those locations, mold growth is inhibited, thereby being able to prevent staining and soiling of wall surfaces, floor surfaces, inside of shoe boxes, shoe surfaces and so forth caused by mold.



[Examples]

[0022] The present invention will be described in more detail by way of the following examples, but the present invention is not limited in any way to these examples.

Example 1 - Confirmation of Fragrant Vapor Antifungal Activity

[0023] The method of Gocho, et al. (Antibacterial and Antifungal Agents, Vol. 19, p. 511-515 (1991)) with a slight variation was carried out to confirm the activity of the fragrant vapor. Physiological saline containing 0.05% Tween-80 was added to a slant on which the test organism grew well, and a spore suspension was produced at  $1 \times 10^6$  CFU (Colony Forming Units)/ml in accordance with ordinary methods. Potato dextrose agar (PDA: Nissui Pharmaceutical) was used for the medium. To begin with, 10 ml of sterile PDA was pipetted into a Petri dish and after it solidified, another 6 ml of PDA containing 1% of above described spore suspension was overlaid and solidfied to prepare the test agar plate. For the fragrance sample, if the sample is in the form of liquid at ordinary temperatures, 20  $\mu$ l of liquid was directly impregnated into a paper disc (Advantech, diameter: 8 mm, thick material), while a 50% by weight ethanol solution was prepared for crystals or powder, and 40  $\mu$ l of that solution was then impregnated into the paper disc. It should be noted that the 40  $\mu$ l of ethanol did not have an effect whatsoever on mold growth.

[0024] The test agar plate was placed inversely, and the paper disc impregnated with the fragrance sample was placed in the center of the cover of the agar plate, followed by incubating for 72 hours at 27°C. Evaluation was made based on the diameter (mm) of the mold growth inhibition zone. When mold growth was inhibited over the entire surface of the agar plate, the diameter of the growth inhibition zone was reported as >85 mm. At locations where mold grew outside the inhibition zone, the diameter of the spore formation inhibition zone and the degree of growth were recorded by comparing with a control. The molds used in the test are shown in Table 1, wherein these molds consist of mold species that are isolated at high frequencies from rooms and bath rooms. In addition, the antifungal activities of the fragrant vapor investigated are shown in Table 2.

Table 1

Abbreviation	Fungi strain name
An-1	Aspergillus niger IFO 9455
Ap-1	Aureobasidium pullulans IFO 6353
Cc-1	Cladosporium cladosporioides IFO 6348

Table 2

Abbreviation of Fungi tested		An-1			Ap-1			Cc-1		
Fragrance name		IZ	SI	Growth	IZ	SI	Growth	IZ	SI	Growth
3-Phenylpropanal		>85		none	>85		none	>85		none
Benzaldehyde		>85		none	>85		none	>85		none
Citral		>85		none	>85		none	>85		none
Citral DEA		>85		sparse	>85		sparse	>85		sparse
Citral DMA		>85		none	>85		none	>85		none
Undecylenealdehyde		>85		none	>85		sparse	>85		sparse
Cuminaldehyde		>85		none	>85		none	>85		none
Furfural		>85		none	>85		none	>85		none
Perillaldehyde		>85		none	>85		none	>85		none
Salicylaldehyde		>85		none	>85		none	>85		none
Trans-2-hexenal		>85		none	>85		none	>85		none
Trans-2-undecenal		>85		none	>85		none	>85		none
Octylaldehyde		20	>85	weak	>85		none	>85		none

## EP 1 214 879 A2

Table 2 (continued)

Abbreviation of Fungi tested		An-1			Ap-1			Cc-1		
5	Nonyl-aldehyde	20		weak	>85		sparse	>85		sparse
	Citronellal	15		weak	>85		sparse	>85		sparse
	Dimethyltetrahydrobenzaldehyde	0		weak	>85		sparse	>85		sparse
10	Cinnamic aldehyde DMA	35	45	weak	65		slight	0		good
	Anisaldehyde	45	60	good	35	40	good	>85		sparse
	Helional	10	35	good	35		weak	0		weak
15	Geraniol	45	65	good	30	45	good	70		weak
	Citronellol	35	55	good	35		weak	35		weak
	9-Decenol	>85		none	65		weak	>85		none
20	Decanol	80		weak	40		slight	60		weak
	Thymol	70	>85	weak	>85		sparse	>85		sparse
	Eugenol	40	60	good	40	60	good	70		weak
25	Hinokitiol	60		good	70		good	0		good
	Perilla alcohol	30	40	good	40		slight	45		slight
	Isoeugenol	30	50	good	35	40	good	45	55	good
30	Methyl salicylate	40		weak	65		weak	0		good
	IZ: Diameter of growth inhibition zone (mm), SI: Diameter of spore formation inhibition zone (mm), Growth: Growth status outside the inhibition zone (Good: Substantially the same as the control, Weak: Somewhat weak growth, Slight: Even weaker growth, Sparse: Low growth density and weak colony growth, None: No growth observed. In addition, a result of "Sparse" for growth of IZ > 85 mm indicates that although an inhibition zone was not formed, test organism growth was weak and sparse.)									

### Example 2 Confirmation of Synergistic Action

**[0025]** An agar plate prepared according to the method of Example 1 was used while using An-1 (*Aspergillus niger*) for the test fungi. To begin with, the amount of each fragrance alone that resulted in a growth inhibition zone of 0 to 30 mm for An-1 was determined, and when that amount was 10  $\mu$ l or less, it was diluted using ethanol, and that 10  $\mu$ l was adjusted to a prescribed amount when impregnated into a paper disc. The test agar plate was placed inversely, and a paper disc impregnated with each fragrance alone and a paper disc simultaneously impregnated with two types of fragrances were placed in the center of the cover, followed by incubating for 72 hours at 27°C and measuring the diameters of the resulting growth inhibition zones. When the combined use of two types of fragrances demonstrated an inhibition zone 20 mm or more larger than the sum of the diameters of the growth inhibition zones of each fragrance alone, the two fragrances were judged to demonstrate synergistic action. When the diameter of the combined growth inhibition zone was 20 mm or more smaller, the two fragrances were judged to demonstrate antagonistic action. When the diameter of the combined growth inhibition zone was intermediate to the above, the two fragrances were judged to demonstrate additive action. A portion of the results were divided into synergistic action, additive action and antagonistic action and shown in Table 3.

Table 3

Sample name	Sample amount	IZ	SI	Growth	Evaluation result
A: Citral	2 $\mu$ l	0	20	weak	Synergistic action
B: Benzaldehyde	10 $\mu$ l	15	>85	slight	
A + B	2 + 10 $\mu$ l	>85	-	none	
A: Dimethyltetrahydrobenzaldehyde	20 $\mu$ l	0	>85	weak	Synergistic action
B: Geraniol	10 $\mu$ l	30	45	weak	
A + B	20 + 10 $\mu$ l	>85	-	none	

Table 3 (continued)

Sample name	Sample amount	IZ	SI	Growth	Evaluation result
A: Octylaldehyde	20 µl	10	35	weak	Additive action
B: 9-Decenol	0.5 µl	0	-	weak	
A + B	20 + 0.5 µl	15	30	weak	
A: Trans-2-undecenal	3 µl	30	>85	slight	Antagonistic action
B: Phenylacetaldehyde	5 µl	25	30	good	
A + B	3 + 5 µl	35	65	weak	

[0026] The meanings of the IZ, SI and growth columns are the same as those shown in Table 2.

[0027] The results obtained in Example 2 are shown in Tables 4 through 7 as combination examples of respective synergistic, additive and antagonistic actions for each fragrant chemical product (a). Those components (b) that demonstrate synergistic action for each fragrant component of component (a) in the present invention are the compound group shown in the "Synergistic Action" column of these tables. It should be noted that in order to slim the table, combinations that have already been listed once are not listed in other columns of the tables. Thus, in the case, for example, the eugenol of Table 7 is component (a), those compounds that demonstrate synergistic action can be found by looking at other fragrance names in the "Synergistic Action" columns of Tables 4 through 7.

Table 4

Fragrance component	Synergistic effect	Additive effect	Antagonistic effect
Citral	Citronellal, Trans-2-hexenal, Octylaldehyde, Trans-2-undecenal, Undecylenealdehyde, Dimethyltetrahydrobenzaldehyde, Benzaldehyde, Salicylaldehyde, Cuminaldehyde, Furfural, Nonylaldehyde, Anisaldehyde, Perillaldehyde, Geraniol, Citronellol, 9-Decenol, Decanol, Thymol, Eugenol, Citral DMA, Citral DEA, Cinnamic aldehyde DMA	3-Phenylpropanal, Phenylacetaldehyde, Methyl salicylate	
Benzaldehyde	Citronellal, Trans-2-hexenal, Octylaldehyde, Trans-2-undecenal, Undecylenealdehyde, Dimethyltetrahydrobenzaldehyde, 3-Phenylpropanal, Salicylaldehyde, Cuminaldehyde, Furfural, Nonylaldehyde, Phenylacetaldehyde, Anisaldehyde, Perillaldehyde, Geraniol, Citronellol, Decanol, Thymol, Eugenol, Citral DMA, Citral DEA, Cinnamic aldehyde DMA, Methyl salicylate		9-Decenol

Table 4 (continued)

Fragrance component	Synergistic effect	Additive effect	Antagonistic effect
Undecylenealdehyde	Citronellal, Octylaldehyde, Trans-2-undecenal, Dimethyltetrahydrobenzaldehyde, Salicylaldehyde, Cuminaldehyde, Nonylaldehyde, Phenylacetaldehyde, Perillaldehyde, Citronellol, 9-Decenol, Decanol, Thymol, Eugenol, Citral DMA, Citral DEA, Cinnamic aldehyde DMA, Methyl salicylate	3-Phenylpropanal, Furfural, Trans-2-hexenal, Anisaldehyde, Geraniol	

Table 5

Fragrance component	Synergistic effect	Additive effect	Antagonistic effect
3-Phenylpropanal	Citronellal, Trans-2-hexenal, Trans-2-undecenal, Dimethyltetrahydrobenzaldehyde, Salicylaldehyde, Citral DMA, Citral DEA	Octylaldehyde, Cuminaldehyde, Furfural, Citronellol, 9-Decenol, Decanol, Thymol, Nonylaldehyde, Anisaldehyde, Perillaldehyde, Cinnamic aldehyde DMA, Eugenol	Geraniol
Furfural	Citronellal, Trans-2-hexenal, Octylaldehyde, Trans-2-undecenal, Cuminaldehyde, Geraniol, Citronellol, Thymol		
Trans-2-hexenal	Citral DEA, Cuminaldehyde, Perillaldehyde, Salicylaldehyde, Thymol, Trans-2-undecenal, Octylaldehyde, Cinnamic aldehyde DMA, Methyl salicylate, Phenylacetaldehyde, Decanol, Eugenol, Anisaldehyde, Geraniol, Dimethyltetrahydrobenzaldehyde, Citronellol	Citral DMA, Nonylaldehyde, 9-Decenol	
Citronellal	Citral DMA, Citral DEA, Cinnamic aldehyde DMA, Methyl salicylate, Cuminaldehyde, Perillaldehyde, Salicylaldehyde, Trans-2-undecenal, Octylaldehyde, Nonylaldehyde, Phenylacetaldehyde, Eugenol, Anisaldehyde, Geraniol, Dimethyltetrahydrobenzaldehyde	Thymol, Decanol, Citronellol	9-Decenol



Table 6

	Fragrance component	Synergistic effect	Additive effect	Antagonism effect
5	Octylaldehyde	Cuminaldehyde, Perillaldehyde, Salicylaldehyde, Trans-2-undecenal, Dimethyltetrahydrobenzaldehyde	Thymol, 9-Decenol, Anisaldehyde, Geraniol, Decanol	Citronellol
10	Trans-2-undecenal	Citral DMA, Citral DEA, Cuminaldehyde, Perillaldehyde, Salicylaldehyde, Decanol, Geraniol, Citronellol	Cinnamic aldehyde DMA, Methyl salicylate	Phenylacetaldehyde, Eugenol
15	Dimethyltetrahydrobenzaldehyde	Cuminaldehyde, Geraniol	Thymol, Decanol	
	Salicylaldehyde	Cuminaldehyde, Thymol, Decanol, Citronellol		
20	Cuminaldehyde	Citral DEA, Geraniol, Cinnamic aldehyde DMA, Phenylacetaldehyde, Decanol, Anisaldehyde, Citronellol	9-Decenol	
25	Geraniol	Nonylaldehyde	Thymol, 9-Decenol	Citronellol, Phenylacetaldehyde, Eugenol

Table 7

30	Perfume component	Synergistic effect	Additive effect	Antagonism effect
	Citronellol	Citral DMA, Citral DEA, Perillaldehyde, Methyl salicylate	Nonylaldehyde, Decanol, Anisaldehyde	Thymol, 9-Decenol, Cinnamic aldehyde DMA, Phenylacetaldehyde, Eugenol
35	9-Decenol	Decanol	Thymol	
40	Decanol		Thymol, Nonylaldehyde, Methyl salicylate, Phenylacetaldehyde, Eugenol	Cinnamic aldehyde DMA
45	Eugenol		Thymol, Methyl salicylate	Phenylacetaldehyde, Anisaldehyde, Cinnamic aldehyde DMA

Example 3 Production of Antifungal Fragrance Composition

[0028] Four types of prepared fragrances (antifungal fragrance compositions) were formulated based on two types of bases (containing neither the components (a) nor (b)) consisting of a citrus-scented base fragrance (citrus base) and floral-scented base fragrance (floral base) according to the results of Examples 1 and 2. The contents of their formulas are shown in Table 8 (values represent parts by weight).

Table 8

	Example 3-1	Comp. ex. 1	Example 3-2	Comp. ex. 2
Citrus base	695	695	0	0
Floral base	0	0	647	647

Table 8 (continued)

	Example 3-1	Comp. ex. 1	Example 3-2	Comp. ex. 2
Component (a)				
Citral	250	0	170	0
Undecylenaldehyde	50	0	30	0
Component (b)				
Benzaldehyde	0	0	150	0
Nonylaldehyde	5	0	0	0
3-Phenylpropanal	0	0	3	0
Etanol (balance)	0	305	0	353

#### Example 4 Antifungal Activity of Compounded Fragrances

**[0029]** The activities of the blended fragrances (antifungal fragrance compositions) prepared in Example 3 were confirmed using the method of Example 1. Those results are shown in Table 9. The amount of blended fragrance added to the paper discs was 20  $\mu$ l each, and only the diameter of the inhibition zone (mm) is indicated for the results.

Table 9

	Fangi tested		
Fragrance compounded	An-1	Ap-1	Cc-1
Example 3-1	>85	>85	>85
Comp. example 1	0	0	0
Example 3-2	>85	>85	>85
Comp. example 2	0	0	0

**[0030]** As can be seen from Table 9, although the blended fragrances of Examples 3-1 and 3-2 containing the component (a) and the component (b) (active ingredients) exhibited potent activity against the three species of bacteria tested, there was no antifungal activity observed for the bases prior to addition of active ingredients.

**[0031]** In addition, separate from the above, the aroma of the blended fragrances were evaluated in the manner described below. Namely, twenty male and female subjects were asked to smell the aromas of Example 3-1, Comparative Example 1, Example 3-2 and Comparative Example 3-2, evaluations were made based on the following evaluation criteria, and the resulting average scores were determined.

Score	Evaluated Contents
4	Retained scent of comparative example
3	Nearly retained scent of comparative example
2	Slight difference from scent of comparative example
1	Large difference from scent of comparative example

**[0032]** The average evaluation score of the prepared fragrance of Example 3-1 was 3.8, while that of the prepared fragrance of Example 3-2 was 3.1. The prepared fragrance of Example 3-1 acquired antifungal activity while retaining the aroma of Comparative Example 1. Although the prepared fragrance of Example 3-2 was observed to exhibit a change in scent, the difference was only slight.

#### Example 5 Preparation of Granular Aromatic Agents

**[0033]** Granular aromatic agents were prepared using the blended fragrances produced in Example 3. Granular types of aromatic agents were produced by using granular calcium silicate (florite, Tokuyama soda) and impregnating with 60% by weight of each blended fragrance of Example 3.

**[0034]** The granular aromatic agent containing the blended fragrance of Example 3-1 was referred to as Example 5-1, the granular aromatic agent containing the fragrance of Comparative Example 1 as Comparative Example 3, the granular aromatic agent containing the blended fragrance of Example 3-2 as Example 5-2, and the granular aromatic

agent containing the fragrance of Comparative Example 2 as Comparative Example 4.

#### Example 6 Antifungal Activity of Granular Aromatic Agents

**[0035]** The granular aromatic agents were evaluated using a vinyl chloride container having an internal volume of 78 liters (inner diameter: 30 cm x 110 cm, stainless steel lining). The test method was as described below. Strips of board cut to a length of 32 cm were grouped into the shape of a cross, four chains were suspended at a location 2 cm from the ends, and four levels of circular metal mesh having a diameter of 25 cm were suspended from the chains. The metal mesh of the first level was positioned 10 cm from the upper edge of the container, and the remaining metal mesh were located at heights of 25 cm, 65 cm and 105 cm from the upper edge of the container. One each of the agar plates of the same bacteria tested used in Example 1 were placed on each metal mesh. The aromatic agents of the present example were used after weighing out so that the amount of active ingredient contained therein was 10 µg/cm<sup>3</sup>. In other words, the total amount of active ingredients shown in Table 8 was calculated to be 780 mg (10 µg/cm<sup>3</sup>) from the scenting rate of the aromatic agents. The weights of the aromatic agents were 4.62 g in Example 5-1 and 3.68 g in Example 5-2. In the comparative examples, aromatic agents were used at the same weights as in the respective examples. Each aromatic agent was wrapped in tissue paper and respectively placed in the center of the metal mesh of the second level. This was done to confirm whether or not the vapor is dispersed upward as well.

**[0036]** After positioning the aromatic agents and agar plates on the metal mesh, they were placed in the container, the opening of the container was wrapped twice and then incubated for 3 days at 27°C. The growth status of the bacteria tested on each agar plate was compared with a control. The results of Example 6 are shown in Table 10.

Table 10

	Example 5-1	Comp. ex. 3	Example 5-2	Comp. ex. 4
First level	No growth	Growth	No growth	Growth
Second level	No growth	Growth	No growth	Growth
Third level	No growth	Growth	No growth	Growth
Fourth level	No growth	Growth	No growth	Growth

**[0037]** The granular aromatic agents of Examples 5-1 and 5-2 completely inhibited growth of the bacteria tested (An-1, Ap-1 and Cc-1) on agar plates placed at the height of the fourth level. In contrast, the bacteria tested on the agar plates placed together with the granular aromatic agents of Comparative Examples 3 and 4 exhibited growth similar to that of the control, and the granular aromatic agents of Comparative Example 3 and 4 were not observed to demonstrate antifungal activity.

#### Example 7 Preparation of Water-Soluble Gel Type Aromatic Agents

**[0038]** Gelatinous aromatic agents were prepared using the blended fragrances produced in Example 3 according to the formulas indicated below.

Formula of Water-Soluble Gel Type Aromatic Agents	
κ-Carrageenan	2.0%
Locust bean gum	0.4%
Polyethylene glycol	3.0%
Calcium chloride	0.4%
Antiseptic (methylparaben)	0.1%
Blended fragrance of Example 3	5.0%
Tween-80	0.5%
Water	88.6%

**[0039]** The gelatinous aromatic agent containing the blended fragrance of Example 3-1 was referred to as Example 7-1, the gelatinous aromatic agent containing the fragrance of Comparative Example 1 as Comparative Example 5, the gelatinous aromatic agent containing the blended fragrance of Example 3-2 as Example 7-2, and the gelatinous aromatic agent containing the fragrance of Comparative Example 2 as Comparative Example 6.

Example 8 Antifungal Activity of Gelatinous Aromatic Agents

[0040] The antifungal activity of the gelatinous aromatic agents was investigated using the same method as Example 6. The weight of the gel used was 51.15 g in Example 7-1 and 44.19 g in Example 7-2. The same weights of gel were also used in each of the comparative examples. The gel was placed in a wide-mouth bottle having an opening diameter of 32 mm, and placed in the center of the metal mesh on the second level in the same manner as Example 6. The antifungal activity of the other gelatinous aromatic agents was investigated in the same manner as Example 6. Those results are shown in Table 11.

Table 11

	Example 7-1	Comp. ex. 5	Example 7-2	Comp. ex. 6
First level	No growth	Growth	No growth	Growth
Second level	No growth	Growth	No growth	Growth
Third level	No growth	Growth	No growth	Growth
Fourth level	No growth	Growth	No growth	Growth

[0041] The gelatinous aromatic agents of Examples 7-1 and 7-2 completely inhibited growth of the bacteria tested (An-1, Ap-1 and Cc-1) on agar plates placed at the height of the fourth level in the same manner as Example 6. In contrast, the gelatinous aromatic agents of Comparative Examples 5 and 6 were not observed to have antifungal activity.

[0042] As has been described in detail above, the antifungal fragrance composition of the present invention contains as its active ingredients (a) at least one type of fragrance component selected from the group consisting of citral, citronellal, trans-2-hexenal, octylaldehyde, trans-2-undecenal, undecylenealdehyde, dimethyltetrahydrobenzaldehyde, benzaldehyde, salicylaldehyde, 3-phenylpropanal, cuminaldehyde, furfural, geraniol, citronellol, 9-decenol, decanol and eugenol, and (b) at least one type of compound that demonstrates synergistic effects in the presence of the above fragrance component selected from the group consisting of aliphatic or aromatic aldehyde, aliphatic alcohol, phenol, acetal and ester, thereby making it possible to reduce the contained amounts of active ingredients to a lower amount than in the case of using each alone.

[0043] Consequently, an antifungal fragrance composition can be produced that retains an arbitrary aroma which has an extremely high degree of safety with respect to the human body. In addition, the blending of these antifungal fragrance compositions with aromatic agents and so forth makes it possible to inhibit the propagation of mold in rooms, closets or shoe boxes, thereby preventing mold from soiling wall surfaces, floor surfaces, shoe surfaces and so forth, or preventing allergies caused by dispersed mold spores.

Claims

1. An antifungal fragrance composition containing as its active ingredients: (a) at least one type of fragrance component selected from the group consisting of an aliphatic or aromatic aldehyde having 5 to 16 carbon atoms, an aliphatic alcohol having 5 to 16 carbon atoms and a phenol; and (b) at least one type of compound demonstrating synergistic effects in the presence of said fragrance component and selected from the group consisting of an aliphatic or aromatic aldehyde, aliphatic alcohol, phenol, acetal and ester; said compound (b) selected being different from said fragrance component (a) selected.
2. The antifungal fragrance composition according to claim 1, wherein said fragrance composition of the component (a) is selected from the group consisting of citral, citronellal, trans-2-hexenal, octylaldehyde, trans-2-undecenal, undecylenealdehyde, dimethyltetrahydrobenzaldehyde, benzaldehyde, salicylaldehyde, 3-phenylpropanal, cuminaldehyde, furfural, geraniol, citronellol, 9-decenol, decanol and eugenol.
3. The antifungal fragrance composition according to claim 1 or 2, wherein the compound demonstrating synergistic effects of the component (b) is citral, citronellal, trans-2-hexenal, octylaldehyde, nonylaldehyde, trans-2-undecenal, undecylenealdehyde, dimethyltetrahydrobenzaldehyde, benzaldehyde, salicylaldehyde, 3-phenylpropanal, cuminaldehyde, furfural, phenylacetaldehyde, anisaldehyde, geraniol, citronellol, 9-decenol, decanol, thymol, eugenol, citraldimethylacetal, citraldiethylacetal, cinnamicaldehydodimethylacetal or methyl salicylate.
4. The antifungal fragrance composition according to any one of claims 1 to 3 that is contained at 1 to 50% by weight in a gel.

**EP 1 214 879 A2**

5. The antifungal fragrance composition according to any one of claims 1 to 3 that is adsorbed or impregnated at 1 to 60% by weight in a powder or granules.
- 5 6. The antifungal fragrance composition according to any one of claims 1 to 3 that is contained at 1 to 30% by weight in a liquid.
7. An antifungal fragrance composition package comprising a gas permeable container which contains the antifungal fragrance composition according to any one of claims 1 to 3, said composition being contained in said container as it is or after having been dissolved in a volatile solvent.
- 10 8. The antifungal fragrance composition package according to claim 7, wherein said gas permeable container is a pouch-shaped object made of a laminate comprising a gas permeable thermoplastic resin on the inside and a gas permeable support on the outside.
- 15 9. The antifungal fragrance composition package according to claim 8, wherein the gas permeable support is a non-woven fabric.
- 20 10. An antifungal fragrance composition package comprising a spray can which contains the antifungal fragrance composition according to any one of claims 1 to 3 and a propellant, said composition being contained in said can as it is or after having been dissolved in a volatile solvent.
- 25 11. A method of inhibiting mold growth comprising: placing the antifungal fragrance composition package according to any one of claims 7 to 9 at an arbitrary location, and dispersing the components (a) and (b) in the antifungal fragrance composition contained in said package.